



## King's Research Portal

DOI:

[10.1017/S0007114518003409](https://doi.org/10.1017/S0007114518003409)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Castro-Acosta, M. L., Sanders, T. A. B., Reidlinger, D. P., Darzi, J., & Hall, W. L. (2018). Adherence to UK dietary guidelines is associated with higher dietary intake of total and specific polyphenols compared with a traditional UK diet: further analysis of data from the Cardiovascular risk REduction Study: Supported by an Integrated Dietary Approach (CRESSIDA) randomised controlled trial. *British Journal of Nutrition*. <https://doi.org/10.1017/S0007114518003409>

### Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### Take down policy

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

## **Title page**

Adherence to UK dietary guidelines is associated with higher dietary intakes of total and specific polyphenols compared to a traditional UK diet: further analysis of data from the CRESSIDA randomised controlled trial.

M.L. Castro-Acosta<sup>1</sup>, T.A.B. Sanders<sup>1</sup>, D.P. Reidlinger<sup>1</sup>, J. Darzi<sup>1</sup>, W.L. Hall<sup>1a</sup>

<sup>1</sup> Diabetes & Nutritional Sciences Division, King's College London, Franklin-Wilkins Building, 150 Stamford Street, London SE1 9NH, UK

<sup>a</sup> Corresponding author: W. L. Hall, email [wendy.hall@kcl.ac.uk](mailto:wendy.hall@kcl.ac.uk)

Department of Nutritional Sciences, School of Life Course Sciences, Faculty of Life Sciences & Medicine, King's College London.

4.108 Franklin Wilkins Building, 150 Stamford St. London

SE1 9NH

Tel: +44 (0)20 7848 4197

**Shortened title:** UK dietary guidelines and polyphenol intakes

**Keywords:** polyphenols, dietary intake, dietary guidelines, randomised controlled trial

## **Abstract**

Adherence to dietary guidelines may result in higher intakes of polyphenols via increased consumption of fruits, vegetables and whole grains. We compared polyphenol dietary intakes and urinary excretion between two intervention groups in the CRESSIDA study: a 12-week, parallel-arm, randomised controlled trial (n=161; 64 M, 97 F; aged 40-70 y). One group adhered to UK dietary guidelines (DG) whereas the other group consumed a representative UK diet (control). We estimated polyphenol dietary intakes, using 4-day food diaries (4-DFD) and food frequency questionnaires (FFQ), and analysed twenty-four hour polyphenol urinary excretion by LC-MS/MS on a subset of participants (n=46 control; n=45 DG). A polyphenol food composition database for 4-DFD analysis was generated using Phenol-Explorer and USDA databases. Total polyphenol intakes by 4-DFD at endpoint (geometric means with 95% CI, adjusted for baseline and gender) were significantly higher in the DG group (1279 mg/d/10 MJ; 1158, 1412) compared to the control group (1084 mg/d/10 MJ; 980, 1197). The greater total polyphenol intakes in the DG group were attributed to higher intakes of anthocyanidins, proanthocyanidins and hydroxycinnamic acids, with the primary food sources being fruits, cereal products, nuts and seeds. FFQ estimates of flavonoid intakes also detected greater intakes in DG compared with the control group. Twenty-four hour urinary excretion showed consistency with 4-DFD in their ability to discriminate between dietary intervention groups for 6 out of 10 selected, individual polyphenols. In conclusion, following UK dietary guidelines increased total polyphenol intakes by approximately 20%, but not all polyphenol subclasses corresponded with this finding.

## 1    **Introduction**

2    Greater consumption of fruit and vegetables and whole grains is associated with reduced risk of chronic  
3    diseases<sup>(1, 2, 3, 4, 5)</sup>. Increased intakes of different components present in these food groups, like fibre<sup>(6)</sup>,  
4    micronutrients<sup>(7)</sup> and polyphenols<sup>(8)</sup>, have been identified as being partly responsible for beneficial  
5    effects. However, results from the UK National Diet and Nutrition Survey (NDNS ) rolling  
6    programme<sup>(9, 10, 11)</sup> reveal that consumption of fruit and vegetables and whole grains is below  
7    recommended intakes in at least 70% of the UK adult population <sup>(9, 10, 11, 12)</sup>. Dietary polyphenols are a  
8    diverse range of phytochemicals containing 1 or more aromatic rings attached to a hydroxyl group. The  
9    term “polyphenol” commonly encompasses phenolic acids, flavonoids, stilbenes and lignans, which are  
10    derived from a wide range of plant foods including fruits, vegetables, and cereals. However, beverages  
11    such as tea, coffee, red wine, and fruit juices represent the main dietary sources<sup>(13)</sup>. Epidemiological  
12    studies have shown a negative relationship between consumption of polyphenols and cardiovascular  
13    disease<sup>(14, 15, 16, 17, 18)</sup>, cancer<sup>(19, 20, 21, 22)</sup> and type 2 diabetes <sup>(18, 23, 24, 25, 26)</sup>. Multiple mechanisms have  
14    been identified that may contribute to any direct causal relationship between dietary polyphenols and  
15    prevention of chronic diseases, including modulating inflammatory pathways, exerting effects on  
16    oxidative signalling and enzyme activity, and regulation of gene expression<sup>(8)</sup>.

17    Previous reports in UK populations, using either 24 h recall<sup>(27, 28, 29, 30, 31)</sup> or food diaries<sup>(27, 28, 29, 30, 31)</sup> ,  
18    suggest that habitual polyphenol intakes are in the range of 800-1600 mg/d, depending on the  
19    population studied. However, the use of different dietary assessment methods, food polyphenol  
20    composition databases, and gaps in polyphenol composition data for certain foods limit the reliability  
21    of current intake estimates in various countries<sup>(32)</sup> . Food diaries may provide a relatively accurate  
22    estimate of polyphenol intake and are more suitable for dietary intervention studies with smaller study  
23    populations, but they reflect short-term intakes rather than habitual consumption patterns, which could  
24    be particularly misleading for seasonally available foods<sup>(33)</sup>. Research groups have designed and  
25    validated FFQs to estimate dietary flavonoid intake in different populations<sup>(34, 35)</sup>, which may provide  
26    more reliable habitual intake estimations for specific populations. However, these remain unavoidably  
27    susceptible to bias due to self-reporting errors, portion size quantification and estimation errors  
28    resulting from the lack of data on polyphenol content in food <sup>(36)</sup>. Whichever dietary assessment  
29    method is selected, the resulting intake data will only be accurate if the polyphenol composition of  
30    foods database is fit for purpose. At present, the most commonly used food polyphenol composition

databases are Phenol-Explorer<sup>(37, 38, 39)</sup>, which provides information on the content of 502 polyphenols (of the 4 classes) in 459 food items including aglycones, glycosides, and esters, and the USDA databases<sup>(40, 41, 42)</sup> which includes 35 flavonoids (aglycones only) in 506 food items, respectively. Urinary excretion of polyphenols has been shown to be a suitable biomarker for the intake of polyphenols<sup>(43, 44)</sup>, fruit and vegetables<sup>(45, 46)</sup>, polyphenol-rich beverages<sup>(47, 48)</sup> and polyphenol-rich food<sup>(49, 50)</sup>. The presence of polyphenol metabolites in urine is closely related to the quantity consumed and overall metabolism in the body. However, different polyphenols can produce common metabolites and so the biomarker selected must reflect the specific intake of the parent polyphenol in question in order to accurately estimate intakes of individual phenolic compounds<sup>(51)</sup>.

Dietary guidelines in the UK are population-based recommendations for maintenance of health and wellbeing, and to reduce risk of chronic diseases. No dietary recommendations exist for polyphenols, although the advantages and disadvantages of this approach have been debated<sup>(52, 53, 54)</sup>. A dietary pattern consistent with current dietary guidelines could reasonably be assumed to be richer in polyphenols than the average UK dietary pattern due to increased intakes of fruit and vegetables and whole grains. However, not all fruits and vegetables are polyphenol-rich<sup>(55)</sup>, and in fact the majority of dietary polyphenols in the UK are provided by tea, coffee, and cocoa intakes<sup>(30)</sup>. The aim of this study was to compare dietary polyphenol intakes in a free-living study population randomised to either following UK dietary guidelines, or consuming a representative, more traditional UK diet. The hypothesis was that adherence to UK dietary guidelines would result in an increase in total dietary polyphenol intakes compared with a diet that was more representative of the UK adult population. The primary outcome variable was total polyphenol intake, adjusted for energy intake, assessed by 4-day food diary. Secondary outcome variables included intakes of classes, subclasses and individual phenolic acids/polyphenols. Data were also compared with FFQ<sup>(15)</sup>, and biomarkers of polyphenol intake (24 h excretion of representative urinary metabolites).

## Methods

### Study design

The Cardiovascular risk REduction Study: Supported by an Integrated Dietary Approach (CRESSIDA) was a 12-week parallel-designed, randomised, controlled trial funded by the Food Standards Agency/Department of Health (UK) (N02047), sponsored by King's College London. This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures

involving human subjects were approved by the St. Thomas' Hospital Research Ethics Committee (10/H0802/24)<sup>(56)</sup>. Written informed consent was obtained from all subjects. This trial was registered at Current Controlled Trial (<http://www.controlled-trials.com/ISRCTN>) as ISRCTN92382106. The primary aim of the original study<sup>(56)</sup> was to assess effects on CVD risk factors when following a diet based on UK dietary guidelines (DG) in comparison to a representative UK diet (control). A sample size of 78/group had 90% power to detect a 4 mm Hg effect of diet on daytime systolic ambulatory blood pressure (alpha 0.05); this sample size was also large enough to detect a 5% change in the ratio of total cholesterol (TC):HDL cholesterol and a 1% unit change in flow-mediated dilatation of the brachial artery with sufficient statistical power. The results of these primary outcomes have been published previously<sup>(56)</sup>.

The data presented here comprises a secondary analysis of dietary intake data from both the DG and control groups at baseline and endpoint of the 12-week dietary intervention period, to determine whether adherence to UK dietary guidelines results in an increase in polyphenol intake.

## **Participants**

Healthy men and women [BMI (in kg/m<sup>2</sup>)  $\geq 18.5$  and  $\leq 35$ ] were recruited from the London area (August 2010–July 2012) by newspaper (London Metro) and electronic advertisement (King's College London e-mail and website). Participants between the age of 40–70 y were chosen because risk increases markedly with age and an upper age limit of 70 y was selected because absolute annual risk of CVD exceeds 2% beyond that age in the majority of people, and a higher proportion are receiving medication on daily basis. A full list of inclusion and exclusion criteria has been reported previously<sup>(56)</sup>. For randomisation a purpose-designed clinical database was used (MedSciNet AB, Stockholm, Sweden), which undertook the minimisation randomisation, balancing the treatment arms for the minimisation variables of gender, age and ethnicity<sup>(56)</sup>. Participants received dietary advice in person at baseline and at week 4, and by phone at week 6 and 8. Dietary assessment included two FFQ and two four-day food diaries (4-DFD) administered at baseline and endpoint and two 24-h recalls administered at weeks 4 and 8. Urine samples were collected at four time points; baseline, week 4, 8 and endpoint.

## **Dietary advice**

Participants randomised to the DG group were advised to increase fruit and vegetables intake to 5 portions/day; whole grains intake to  $>50\%$  of cereal intake; to consume 2 portions of fish per week (1 of which should be oily); to replace full-fat with reduced fat dairy products; to replace fats rich in

91 saturated fatty acids with spreads/oils low in saturated fatty acids and high in monounsaturated fatty  
92 acids; to restrict salt intake to <6 g/d (<100 mmol/d) and to reduce intake of free sugars. The control  
93 diet comprised a nutritionally balanced, traditional UK diet, formulated with familiar foods (full cream  
94 milk, cheese, butter, meat and meat products, non-wholegrain cereals), reflecting typical UK intakes of  
95 fruit and vegetables (3 portions/d), with a higher content of saturated fatty acids (14% energy),  
96 unrestricted intakes of salt and sugar, and low intakes of oily fish. The study included provision of  
97 spread, oil, whole grain pasta, rice and cereal bars, minimally processed wholegrain breakfast cereal  
98 (oats, muesli, etc.), almonds and macadamia nuts and tinned oily fish to the intervention group and  
99 spread, oil and refined cereals as pantry items to the control group. Participants of both groups were  
100 instructed to refrain from taking nutritional supplements during the study<sup>(57)</sup>.

#### 101 **Four-day food diary (4-DFD)**

102 Dietary intake of polyphenols was quantified from 4-DFD, at baseline and endpoint of the 12-wk  
103 intervention. A polyphenol food composition database was generated using Phenol-Explorer<sup>(37, 38, 39)</sup>  
104 and USDA databases<sup>(40, 41, 42)</sup>, and extended using polyphenol retention factors and recipes provided by  
105 participants. If a recipe was not provided by a participant, a standard recipe was obtained from either  
106 the UK food tables<sup>(58)</sup>, a UK food industry recipe book<sup>(59)</sup>, the BBC Good Food website<sup>(60)</sup> or additional  
107 websites specialised in UK recipes and other countries' typical cuisines. In total 118 recipes were  
108 obtained from participants, 93 from UK food tables, 20 from a UK food industry recipe book, 78 from  
109 the BBC Good Food website and 94 from other websites. Polyphenol intake was estimated for four  
110 classes of polyphenols: flavonoids, phenolic acids, lignans and stilbenes. An additional group of  
111 polyphenols were included under the name "other polyphenols"; since Phenol-Explorer grouped a  
112 series of compounds including alkylmethoxyphenols, alkylphenols, curcuminoids, furanocoumarins  
113 and tyrosols. The flavonoid intake was analysed in eight subclasses: anthocyanins, dihydrochalcones,  
114 flavanols (flavan-3-ols monomers and theaflavins), proanthocyanidins, flavanones, flavones, flavonols  
115 and isoflavones, for each subclass a range of 2 to 11 of the most representative individual compounds  
116 were selected. For phenolic acids subclass 2 groups were analysed: hydroxybenzoic acids and  
117 hydroxycinnamic acids, for each group a range of 4 to 6 of the most representative individual  
118 compounds were selected. For lignans subclass, 4 individual compounds were analysed: pinoresinol,  
119 lariciresinol, secoisolariciresinol and matairesinol. For stilbenes subclass, resveratrol was selected and  
120 analysed. A total of 52 individual polyphenols were analysed and 1141 food items were included in the  
121 final database.

## **Food frequency questionnaires (FFQ)**

Dietary intake of flavonoids was quantified from FFQ (EPIC-Norfolk FFQ v.6)<sup>(61)</sup> at baseline and endpoint of the 12-wk intervention. Flavonoid intake was estimated from 6 subclasses and most representative compounds for each: flavanones (hesperetin, naringenin, eriodictyol), anthocyanins (cyanidin, delphinidin, malvidin, pelargonidin, peonidin, petunidin), flavones (apigenin, luteolin), flavonols (quercetin, kaempferol, myricetin, isorhamnetin), flavan-3-ols (catechin, epicatechin, gallocatechin, epigallocatechin, epicatechin-3-gallate, epigallocatechin-3-gallate) and polymers (theaflavins, thearubigins and proanthocyanidins). In total 32 individual flavonoids were analysed using a database created and kindly provided by Dr Amy Jennings and Prof Aedin Cassidy at University of East Anglia<sup>(15)</sup> and modified for the requirements of the FFQ analysis, so that the flavonoid content of 130 food items were analysed. FFQ registers the frequency of consumption of specific foods, of standard portion size, in a month. There were nine answer options which varied from “never or less than once in a month” to “six or more per day”. To calculate the total intake of subclasses of flavonoids, frequencies were converted to daily portions and multiplied by flavonoid content in each specific food item; the results were summed for each subclass and for each participant.

## **Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS)**

Twenty-four hour urine collections were made as previously described<sup>(56)</sup>, using boric acid as preservative<sup>(62)</sup> and aliquots stored at -80 °C until analysis. Completeness of urine collection was measured using recovery of para-aminobenzoic acid according to a standard protocol<sup>(63)</sup>. LC-MS/MS analysis was only performed on endpoint urine samples, collected at week 12 of the intervention to coincide with dietary assessment by 4-DFD. A previously published protocol for polyphenol analysis in urine<sup>(47)</sup> was adapted and developed for the quantification of ten aglycone metabolites of polyphenols. A subsample of the CRESSIDA study was selected for the analysis; the selection was based on the fruit and vegetables (F&V) intake reported in the endpoint FFQ, subsample included participants in the control group reporting the lowest intakes of F&V, and participants in the intervention group reporting the highest intakes of F&V were analysed. Quantification of endpoint samples allowed the comparison between groups at the end of the dietary intervention; the aim was to detect greater amounts of polyphenols excreted in urine by participants who reported consuming the most polyphenols by FFQ compared to participants who reported consuming the least, and to determine agreement with food diary data. One or two of the most representative aglycones of different polyphenol subclasses were selected for quantification. Ten phenolic compounds were analysed; phloretin (dihydrochalcones),



153 epicatechin (flavan-3-ols), hesperetin and eriodictyol (flavanones), luteolin (flavones), quercetin  
154 (flavonols), daidzein (isoflavones), gallic and vanillic acid (phenolic acids) and enterolactone as a  
155 product of microbial metabolism of lignans in colon. The phenolic compounds selected for  
156 quantification have been suggested as suitable biomarkers of polyphenol intake<sup>(55)</sup> with a high recovery  
157 and good correlation with fruit and vegetable intakes when estimated by FFQ, 24-h recalls or food  
158 diaries<sup>(50)</sup>. The major food sources of the ten phenolic compounds are fruit, vegetables and wholegrain  
159 products, or represent the main metabolites of parent glucosides present in the previously mentioned  
160 foods.

161 Purified standards phloretin, epicatechin, eriodictyol, hesperetin, luteolin, daidzein, gallic and vanillic  
162 acids were obtained from Santa Cruz Biotechnology, Germany and quercetin, enterolactone and  
163 catechin-2, 3, 4-<sup>13</sup>C<sub>3</sub> from Sigma-Aldrich, UK. Stock solutions were prepared for all the purified  
164 phenolic compounds, two milligrams of each of the ten purified compounds were diluted in one  
165 millilitre of methanol or dimethyl sulfoxide according to supplier specifications and one milligram of  
166 Catechin-2, 3, 4-<sup>13</sup>C<sub>3</sub> (internal standard) in one millilitre of methanol. Stock solutions were stored at -40  
167 °C or -80 °C (internal standard). Enzyme  $\beta$ -glucuronidase/sulfatase (0.05 g) type H-5 from *Helix*  
168 *Pomatia* (Sigma-Aldrich, UK) was diluted in one millilitre of 0.2% sodium chloride solution to create a  
169 working solution of 50,000 units per millilitre; the enzyme solution was stored at -40 °C. Urine  
170 samples (250  $\mu$ l) were processed for hydrolysis of glucuronide and sulphated metabolites using a  
171 modified version of Ito *et al.* <sup>(47)</sup>. Urine samples were acidified with acetic acid (20  $\mu$ l, 0.58 M) and  
172 incubated with 1300 units of  $\beta$ -glucuronidase/sulfatase and 300 ng of internal standard at 37 °C, 120  
173 rpm for 1.5 h. A liquid-liquid extraction with ethyl acetate was performed twice (400  $\mu$ l and 300  $\mu$ l).  
174 The two organic layers were pooled and evaporated to dryness under N<sub>2</sub> then stored at -80 °C for 1-4  
175 days until their reconstitution with 100  $\mu$ l of 40% methanol immediately before injection. Analysis of  
176 urine samples was performed on a HPLC system Hewlett-Packard series 1100 binary pump, coupled to  
177 a triple quadrupole mass spectrometer, Micromass Quattro LC (Micromass, Limited), operating in  
178 negative electrospray ionization (ES-) mode, equipped with a Zorbax SB-C18 column (2.1 x 50mm,  
179 3.5mm, Agilent). Ionization and fragmentation were optimized for each polyphenol by direct infusion  
180 of a standard solution and specific values for collision energy parameters were identified for each  
181 polyphenol. Peak for each polyphenol identity was established by the parent and daughter ion pair peak  
182 and retention time. Solvents A (water with 0.1% v/v formic acid) and B (acetonitrile with 0.1% v/v  
183 formic acid) were run in a 95/5% proportion at a flow rate of 0.2 ml/min with the following gradient:

184 0–1 min, 5% solvent B; 1–4 min, increase solvent B from 5% to 10%; 4–5 min, increase solvent B  
185 from 10% to 90%, 5–5.2 min, decrease solvent B from 90% to 5%, 5.2–15 min isocratic for 9.8 min.  
186 Peak areas were plotted against the internal standard response. A good linearity ( $r^2$  0.970–0.990) was  
187 observed for all the polyphenols quantified except for luteolin ( $r^2$  = 0.922). Samples were run in the  
188 same batch and chromatograms were processed automatically by MassLynx Mass Spectrometry  
189 Software (MASSLYNX™ version 3.5) using the same processing integrate parameters, peak-to-peak  
190 amplitude and peak detection. Intra-assay coefficients of variability were phloretin 7.5%, epicatechin  
191 6%, hesperetin 21.9%, eriodictyol 9.6%, luteolin 9.8%, quercetin 10.8%, daidzein 3.1%, gallic acid  
192 21.4%, vanillic acid 14.4% and enterolactone 5.1%. Final quantities of phenolics compounds were  
193 estimated after adjustment by urinary volume. Calibration curves were prepared by spiking HPLC-  
194 grade water with 2.5, 5, 12.5, 25, 50, 100, 150, 200 and 250 µl of mixed polyphenols working solution  
195 and 100 µl of internal standard working solution, the calibration curve range was 5–500 ng/ml. Spiked  
196 HPLC-grade water samples were treated with enzyme and extracted with ethyl acetate as were the  
197 urine samples, spiked samples were injected in duplicate at each concentration level. Limits of  
198 quantification were established using the spiked samples for calibrations curves; minimum detected  
199 values were 5 ng/ml for phloretin, epicatechin, eriodictyol, hesperetin, luteolin, daidzein, gallic acid  
200 and vanillic acid and 10 ng/ml for quercetin and enterolactone.

## 201 **Statistical analysis**

202 Statistical analysis was performed using Statistical Package for the Social Sciences (SPSS) v.21. Four-  
203 day food diary intake data<sup>(64)</sup> were expressed as weight of intake (g) per day per 10 MJ energy intake,  
204 in order to adjust for variability in total food intake between individuals over the 4 days, whereas FFQ  
205 data were expressed simply as weight of intake (g) per day. Normality of data distribution was  
206 evaluated visually by inspection of histograms and normal Q-Q plots. Non-normally distributed  
207 variables were natural log transformed prior to statistical analysis by parametric methods, or analysed  
208 by non-parametric methods where log transformation failed to yield a normal distribution. Independent  
209 sample t-tests were conducted to compare groups at baseline in order to verify that the polyphenol  
210 intakes of each treatment group were similar before the dietary intervention. Where data were normally  
211 distributed, one-way analysis of variance with gender as a fixed factor and baseline value as covariate  
212 (ANCOVA) was conducted to find differences between groups at endpoint. Where data could not be  
213 transformed to a normal distribution, a Mann-Whitney U test was conducted to find differences

214 between groups at endpoint. Correlation analyses on urinary and dietary polyphenols were conducted  
215 by two-tailed Spearman's correlations.

## 216 **Results**

217 A total of 165 healthy men and women aged 40–70 years were recruited, and 162 completed the  
218 CRESSIDA study (64 M, 97 F) (**Figure 1**). Dietary intake of polyphenols was quantified from 322 x 4-  
219 DFD (n=161) and from 322 FFQ (n=161) (EPIC-Norfolk FFQ v.6)<sup>(61)</sup> at baseline and endpoint of the  
220 12-wk intervention, one participant randomised to the DG group did not complete the collection of  
221 food diaries therefore was eliminated from the analysis. Urinary excretion of polyphenols was analysed  
222 in a subsample (n=91) of the CRESSIDA study, 45 samples from participants in the control group  
223 reporting the lowest intakes of F&V, and 46 samples from participants in the intervention group  
224 reporting the highest intakes of F&V.

225 Characteristics of the whole CRESSIDA study population at baseline are shown in **Table 1**. Mean BMI  
226 was significantly higher in the control group. Intakes of wholegrain cereals, fruits and vegetables and  
227 dietary fibre were higher in the study population at baseline than those reported in a larger,  
228 representative UK population sample <sup>(11, 56)</sup>. As reported previously, at endpoint urinary potassium  
229 excretion was 9 mmol/d greater in the DG group indicating higher consumption of fruit and vegetables.  
230 Reported wholegrain cereal intake was 81 g/d at endpoint in the DG group (mainly wheat, oats and  
231 rice) compared with 32 g/d in the control group, as confirmed by higher plasma alkylresorcinol  
232 concentrations, reflecting intakes of whole grains mainly from wheat, barley, and rye <sup>(56)</sup>.

### 233 **Four-day food diary**

234 Median total polyphenol intake at baseline in the whole study population (with lower and upper limits  
235 of IQR) was 1183 mg/d (745, 1613), or 1282 mg/d per 10 MJ energy intake (896, 1838). As shown in  
236 **Table 2**, baseline intakes of total polyphenols were significantly higher in the DG group compared to  
237 the control group due to greater intakes of total flavonoids, the main contributor being total  
238 proanthocyanidins.

239 At endpoint, total polyphenol intakes per 10 MJ energy intake, adjusted for gender and baseline  
240 polyphenol intake, were higher in the DG group than in the control group (**Table 2**). Analysis of  
241 intakes of polyphenol classes at endpoint (unadjusted for gender and baseline as non-parametric  
242 statistical analysis was necessary) demonstrated that the DG group had higher intakes of total

243 flavonoids (although also higher in DG at baseline), total lignans, phenolic acids, stilbenes, and “other  
244 polyphenols” compared to the control group. For the major flavonoid subclasses, the DG group  
245 reported higher intakes of anthocyanidins, proanthocyanidins (although these were also higher at  
246 baseline), and isoflavones, but not dihydrochalcones, flavanols, flavones or flavonols. For the major  
247 phenolic acids subclasses, the DG group reported higher intakes of hydroxycinnamic acids, but the  
248 trend for higher intakes of hydroxybenzoic acids was not statistically significant.

249 For individual (poly)phenols, there were no differences between groups at baseline, but at endpoint the  
250 group following the DG diet reported higher intakes of a number of anthocyanidins including cyanidin,  
251 malvidin, peonidin and petunidin ( $P<0.05$ ) compared to the control group. Furthermore, intakes of  
252 individual flavanones (naringenin, eriodictyol, and hesperetin,  $P<0.05$ ), a flavone (luteolin,  $P<0.05$ ),  
253 and isoflavones (daidzein and genistein,  $P<0.01$ ) were higher in the DG group compared to the control  
254 group at endpoint. With regards to lignans, phenolic acids and stilbenes classes, there were higher  
255 intakes of secoisolariciresinol ( $P<0.05$ ), matairesinol ( $P<0.001$ ), protocatechuic acid ( $P<0.01$ ), vanillic  
256 acid ( $P<0.001$ ), 4-hydroxybenzoic acid ( $P<0.001$ ), syringic acid ( $P<0.001$ ), p-coumaric acid ( $P<0.001$ )  
257 and ferulic acid ( $P<0.05$ ) at endpoint in the DG group compared to the control group. There were no  
258 differences between groups in individual flavonols (including kaempferol, quercetin, myricetin and  
259 isorhamnetin) nor flavanols (including individual catechins and theaflavins).

260 The main sources of total polyphenols at endpoint in both groups were tea, coffee & wine (control;  
261 61%, DG; 59%), fruits (control; 10%, DG; 17%) and fruit juices (control; 8%, DG; 4%) (**Figure 2**).  
262 Main sources of flavonoids in control and DG groups were tea, coffee & wine (control; 55%, DG;  
263 55%), fruits (control; 14%, DG; 21%) and fruit juices (control; 10%, DG; 5%). Main sources of the  
264 flavanols subclass were tea, coffee & wine (control; 64%, DG; 65%), fruits (control; 13%, DG; 19%)  
265 and chocolates (control; 11%, DG; 4%). There was an increase in the percentage of polyphenols  
266 sourced by fruits, cereal products (breakfast cereal and cereal bar) and nuts & seeds (see **Figure 3** for  
267 further details) in the group following the DG diet compared to baseline. There was a decrease in  
268 percentage of polyphenols derived from less recommended food options as fruit juices, chocolates and  
269 biscuits (**Figure 3**).

## 270 Food frequency questionnaire

271 There were no differences in total flavonoids and flavonoid subclass estimated intakes between groups  
272 at baseline. At endpoint the DG group reported higher intakes of total flavonoids, total anthocyanidins,

total flavones, total flavanols, and total proanthocyanidins (**Table 3**). Flavanone and flavonol intakes did not differ between groups at endpoint. Differences between individual polyphenol components revealed higher reported intakes of cyanidin, delphinidin, petunidin and peonidin ( $P<0.001$ ), malvidin and pelargonidin ( $P<0.005$ ), catechin ( $P<0.01$ ), and epicatechin-3-O-gallate, epigallocatechin-3-O-gallate, myricetin and luteolin ( $P<0.05$ ) in the DG group (data not shown) compared to control.

### **Urinary polyphenol excretion**

The ten aglycones quantified in urine were either metabolites of parent glucuronide/sulphated compounds formed in human tissues, or metabolites produced by the gut microbiota, e.g. enterolactone from lignans present in fibre-rich food. At endpoint there was a greater 24 h urinary excretion of phloretin, eriodictyol, hesperetin, luteolin, quercetin, gallic acid, vanillic acid and enterolactone in the DG group compared to the control group (**Table 4**), and a non-significant tendency towards a similar response for daidzein excretion. Daidzein was not detected in 37 participants of the 91 subsample, which could contribute to the lack of significance in the results given that values observed are considerably higher in the intervention group. The only polyphenol that was excreted in similar amounts in both dietary groups was epicatechin.

### **Discussion**

The aim of this secondary analysis of dietary intake data from the CRESSIDA trial was to investigate whether following advice to adhere to UK dietary guidelines increased polyphenol intakes. Baseline polyphenol/flavonoid intakes from 4-DFD and FFQ were in alignment with previous reports in UK populations using 24 h recall<sup>(27, 28, 29, 30, 31, 65, 66, 67)</sup> or FFQ<sup>(15)</sup>. The hypothesis was supported by the finding that total estimated polyphenol intakes were approximately 200 mg per 10 MJ energy intake higher at endpoint in the DG group compared to the control group. However, estimates of total polyphenol intakes are only as accurate as the polyphenol composition data, and rely on the sum of the components, which may lead to underestimates if complete composition profile data are unavailable for individual foods. Therefore, total polyphenol intake data should be interpreted in the context of the individual polyphenol intakes. Estimated dietary intakes by 4-DFD showed that the DG group had higher intakes of individual polyphenols/phenolic acids where main food sources were fruits and vegetables (cyanidin, malvidin, peonidin, petunidin, protocatechuic acid, naringenin, eriodictyol, hesperetin and syringic acid), nuts and seeds (secoisolaricresinol and matairesinol) and soy products (daidzein, genistein and other isoflavonoids) compared to the control group. In agreement with the 4-

303 DFD results, FFQ estimates of total flavonoids, proanthocyanidins and anthocyanidins were higher in  
304 the DG group compared to the control group. Ten representative aglycone polyphenols and phenolic  
305 acids (from glucuronidated and sulphated metabolites) were measured in 24 h urine samples collected  
306 concurrently with 4-DFD as objective biomarkers of (poly)phenol intakes, in order to determine the  
307 accuracy of the dietary intake estimates. The ability to discriminate between the dietary group  
308 subsamples was consistent with 4-DFD, except for epicatechin, daidzein and quercetin where there  
309 were no significant differences in urinary excretion in contrast with higher estimated intakes in the DG  
310 group by 4-DFD. Both the 4-DFD and urinary biomarker methods agreed that intakes of  
311 phlorizin/phloretin, eriodictyol, hesperidin, luteolin, vanillic acid and lignans/enterolactone were  
312 greater in the DG group.

313 The ability to discriminate between dietary intervention groups supports the utility of 4-DFD and  
314 urinary metabolites for assessment of dihydrochalcones, flavanones, flavones, hydroxybenzoic acids  
315 and lignan intakes over short-term periods. Differing findings may be due to incomplete phenolic  
316 composition data, the well-known inaccuracies of self-reported dietary assessment methodology<sup>(68, 69)</sup>,  
317 and the inter-individual variability in absorption and metabolism<sup>(8, 70)</sup> or incomplete deconjugation of  
318 sulphated/glucuronidated metabolites<sup>(71)</sup> reflected in the urinary biomarkers. Urine polyphenol  
319 concentrations may also be the sum product of endogenous metabolism and dietary intakes. For  
320 example, gallic acid is a common metabolite of black tea theaflavins and thearubigins<sup>(72, 73)</sup>, therefore  
321 its urinary excretion could reflect the consumption of such beverage combined with foods providing  
322 gallic acid *per se*. Vanillic acid can be found in high concentrations in foods consumed in small  
323 quantities such as herbs; also in moderate concentration in dates, olives and cranberries and in low  
324 concentrations in foods more frequently consumed like oats and rice<sup>(39)</sup>. It is also a common metabolite  
325 of anthocyanins<sup>(74, 75)</sup>, and so variability in urinary vanillic acid excretion could reflect varying intakes  
326 of both vanillic acid and anthocyanin-rich foods, therefore it is likely that the increased urinary  
327 excretion at endpoint mainly represents intake of foods rich in anthocyanins rather than vanillic acid.  
328 Vanillic acid is also reported to be present in urine after consumption of black tea<sup>(76)</sup> and dark  
329 chocolate<sup>(77)</sup>; however, intakes of black tea and dark chocolate were not higher in the DG group at  
330 endpoint, and therefore are unlikely to account for the increased urinary excretion and reported intakes  
331 in the 4-DFD. Enterolactone is the main colonic metabolite of lignans, which are phytoestrogenic  
332 compounds present in high concentrations in vegetables, cereals and grain products, seeds, nuts and  
333 berries and other fruits<sup>(65, 78)</sup>. These urinary excretion data are in agreement with previous  
334 epidemiological studies<sup>(79)</sup>, supplementation studies where known doses of polyphenols were

335 administrated<sup>(44, 80)</sup>, in RCTs aiming to increase flavonoid-rich fruit and vegetables intake<sup>(81)</sup> and with  
336 population under free-living conditions<sup>(43)</sup>. Polyphenol urinary excretion has shown to be an useful  
337 biomarker of fruit and vegetables intakes, capable of detecting even small changes in studies under  
338 controlled diets<sup>(45, 82, 83)</sup>.

339 Following dietary guidelines did lead to increases in specific polyphenols that have been associated  
340 with health benefits (e.g. anthocyanins, flavanones, isoflavones, phenolic acids and lignans<sup>(84, 85, 86, 87)</sup>.  
341 This suggests that consuming at least 5 portions F&V per day and >50 % of cereal intake as wholegrain  
342 will lead to an enrichment in dietary polyphenol profiles above and beyond the large amounts of  
343 polyphenols supplied by beverages. In particular, estimated anthocyanidin intakes were augmented by  
344 following dietary guidelines which may confer a reduction in risk of T2D<sup>(25)</sup>, and acute reductions in  
345 postprandial glycaemia<sup>(88, 89, 90, 91)</sup>. The dietary intake results reported here are consistent with the theory  
346 that greater intakes of specific polyphenols may contribute to the protective effects of a diet rich in  
347 fruits and vegetables and wholegrain cereals, alongside greater dietary intakes of nutrients and other  
348 non-nutrient bioactives, e.g. potassium, vitamin C, soluble fibre, carotenoids and glucosinolates.  
349 However, our results confirm<sup>(29, 30, 31)</sup> that food groups that are recommended by UK government as  
350 part of the Eatwell Guide – fruits and vegetables, wholegrain cereals, beans, pulses, nuts and seeds -  
351 contribute a much smaller proportion of daily polyphenol intakes in a UK population compared to  
352 commonly consumed beverages, even when that population is following dietary guidelines. Analysis of  
353 dietary data showed no group differences in (poly)phenol intakes (by 4-DFD) where main food sources  
354 were not targets for UK dietary guidelines such as tea (catechins, theaflavins, proanthocyanidins,  
355 quercetin) and wine (stilbenes, proanthocyanidins). However, the sources of these polyphenols may  
356 have differed between groups; for example, the DG group reported lower intakes of stilbenes from  
357 wine and increased intakes of stilbenes from fruit.

358         The relative impacts of total and specific (poly)phenol intakes on risk of CVD and type 2  
359 diabetes requires investigation. Clearly there is only limited scope for increasing total polyphenol  
360 intake by following dietary guidelines – an approximate increase of 200 mg per day per 10 MJ energy  
361 intake. The extent to which this is relevant to cardio-metabolic health is currently unclear due to the  
362 lack of robust evidence for the relative impact of different polyphenol subclasses on risk factors for  
363 CVD and type 2 diabetes. Although the evidence is currently insufficient, it is plausible to hypothesize  
364 that in the future there may be consensus that tea and coffee (poly)phenols<sup>(92, 93)</sup> may have some of the  
365 strongest cardio-metabolic protective effects amongst all dietary phenolics, so that due to the large

amounts of tea and coffee consumed at a population level, the gain to health in consuming at least 5 portions/day of fruits and vegetables and choosing wholegrain sources of starchy carbohydrates is likely to be due to other nutritional factors. Conversely, future advances in nutritional science may one day demonstrate that certain polyphenols specific to fruits, vegetables or wholegrain cereals have particularly potent bioactivity in preventing inflammation, atherosclerosis or insulin resistance, which would bring about a departure from the prevailing approach of considering all plant chemicals that contain a similar chemical structure in the same way. There has been some debate around whether establishing dietary reference intakes (DRI) for polyphenols could be beneficial; with suggested approaches including recommendations for polyphenol-rich F&V (5-a-day)<sup>(52)</sup>, establishing a specific daily dose for a given effect<sup>(54)</sup> or establishing values to improve health or prevent disease risk in different life stages<sup>(53)</sup>. Further research is required to provide robust interventional evidence for any refinement to current “5-a-day” guidelines, since the relative health impact of a multitude of other non-nutrient bioactives and the complex array of plant cell wall polysaccharides, resistant starch and oligosaccharides (collectively known as fibre) contained in low-polyphenol plant-based foods is only partially understood. Furthermore, in the light of the fact that NDNS survey data suggest that the UK population are not meeting current recommendations for F&V then it would be futile, and possibly counter-productive, to add further complexity to existing public health dietary advice.

The validation of adequate biomarkers of dietary intake, of exposure to dietary components and of compliance to dietary interventions has been a focus of investigation for many years<sup>(94)</sup> and the identification of polyphenol intake biomarkers is no exception<sup>(83, 95)</sup>. Assessment of compliance in dietary intervention studies with fruits and vegetables has been done by validated biomarkers<sup>(96)</sup> such as vitamin C, carotenoids and potassium, however polyphenol excretion in urine may be used to assess flavonoid-rich dietary interventions<sup>(81)</sup> or specific fruit and vegetable intakes, for example phloretin for apple intake and naringenin and hesperetin for citrus fruit<sup>(95, 97)</sup>. Given the importance of accurate assessment of polyphenol intake in order to link it with beneficial effects on health<sup>(98)</sup>, the ability to detect the amount consumed represents a challenge in epidemiological and clinical studies.

The bespoke food composition table developed for the 4-DFD polyphenol analysis in the present study included the estimation of 52 individual compounds and contained 1141 foods representative of an average UK diet as well as some of the most commonly eaten food from non-traditional UK cuisine as Turkish, Indian, Italian, Chinese, Japanese, and others. This food composition table could be used as a basis for future studies analysing diet patterns in the UK population, comparing



polyphenols intake and/or linking intake with health benefits, although it would require a significant staff cost commitment to maintain and update with new recipes and new versions of Phenol Explorer over time. Although an onerous process, the creation of a bespoke food composition table for polyphenol analysis in a specific population seems to be the most effective way to obtain an accurate estimation of polyphenol intake. Although the FFQ is a valid and reliable dietary assessment tool for estimation of habitual intake in populations, it has limitations in the way that it can collect information, since it is based on uniform estimated portion sizes and is semi-quantitative. Dietary intake of total flavonoid intake, six flavonoid subclasses and 32 individual flavonoids (aglycones) were estimated by FFQ (EPIC-Norfolk) using methodology previously applied in large cohort studies<sup>(15, 17, 23, 25, 26, 99)</sup>. The EPIC-Norfolk FFQ, although validated for whole foods intake<sup>(100)</sup>, has not been validated for flavonoid intake. Methodological barriers of this approach in regard to flavonoid estimation are particularly limiting in smaller study cohorts: some foods are grouped incongruously with regards to their flavonoid composition such as “strawberries, raspberries and kiwi”, or “peaches, plums and apricots” and “wine” (no separation of red, white and rosé). Some important flavonoid sources are missing such as blueberries, soy beverages, and lemon, among others. The inadequacy of this method could lead to under- or overestimation of flavonoid intakes. Studies have used food diaries for estimation of flavanol intake in European populations<sup>(29)</sup>, isoflavone and lignan intakes<sup>(66)</sup> and 20 subclasses of polyphenols<sup>(31)</sup> in a UK population, each one creating their own food composition table for polyphenol analysis, however they have not included objective biomarkers for corroborating agreement between methods. More specialised questionnaires for intake estimation have been developed for individual compounds like quercetin and naringenin<sup>(101)</sup>, flavonoid subclasses such as flavonols and flavones in a Chinese population<sup>(102)</sup>, complete flavonoid class in Australian population<sup>(34)</sup> and Flemish population<sup>(103)</sup>. In summary, although the FFQ results reported here were in broad agreement with the 4-DFD regarding the discernment of differences in total flavonoid intakes in a dietary intervention study, future research in this area could include the development of a specific FFQ, validated in a UK population, for estimation of phenolic intakes including all classes and main subclasses as were estimated by 4-DFD in this study.

Limitations in the present study include the hydrolysis of urinary polyphenols by glucuronidase/sulphatase enzymes prior to analysis, which could compromise the stability of polyphenols that are to be quantified<sup>(104, 105)</sup>. Glucuronidase/sulphatase enzymes may fail to completely hydrolyse all sulphated and methylated-sulphated metabolites, which may have been compounded by the addition of boric acid as a preservative to prevent yeast growth, and therefore urinary excretion of

429 the corresponding aglycone could be underestimated<sup>(71)</sup>. Furthermore, our approach required  
430 preselection of polyphenols to be analysed which means that many other polyphenols present in 24 h  
431 urine that may have also functioned as biomarkers of intake were not detected and quantified. Although  
432 reported anthocyanidin intakes were greater in the DG group, we did not attempt to analyse  
433 anthocyanins in urine as they have low recovery as parent compounds and only weak correlations with  
434 dose ingested, necessitating the use of stable isotopes to accurately measure the excretion of  
435 anthocyanin metabolites<sup>(44)</sup>. The 4-DFD proved to be an accurate approach for estimate of short-term  
436 polyphenol intake, however, serious consideration must be made of the potential researcher burden and  
437 cost if using these methods in larger scale studies, particularly where a simple indicator of compliance  
438 to dietary advice is required.

439 Strengths of our study include the fact that the data collected in this study enabled a comparison of two  
440 dietary assessment methods with objective biomarkers of (poly)phenol dietary intake collected  
441 concurrently towards the end of the dietary intervention period. To date, this 3-way comparison has  
442 only been reported previously in a cross-sectional<sup>(106)</sup> study using a crude estimate of total urinary  
443 polyphenol excretion by the Folin-Ciocalteu assay. To the best of our knowledge this is the first study  
444 to use the 3 way comparison on data obtained from an RCT allowing comparison of change from  
445 baseline for individuals advised to consume a diet consistent with dietary guidelines.

446 In conclusion, participants following advice to adhere to UK dietary guidelines consumed a greater  
447 amount of total polyphenols than a control group consuming a representative UK diet. In particular,  
448 intakes of individual polyphenols that are mainly sourced from fruits and vegetables, nuts, seeds and  
449 soy products were increased in the DG group. The results of this study advance the field of polyphenol  
450 research as they demonstrate that a diet consistent with dietary guidelines is also moderately richer in  
451 polyphenols compared to a typical UK dietary pattern, despite the relatively high baseline intakes  
452 derived from commonly consumed beverages such as tea and coffee. In addition to increased intakes of  
453 dietary fibre and certain micronutrients, replacement of saturated fatty acids with unsaturated fatty  
454 acids, and lower salt and free sugars, following dietary guidelines will also increase polyphenol intakes  
455 which may contribute to the overall reduction in risk factors for cardiometabolic diseases.

## **Acknowledgments**

We thank Anna Caldwell (CEMS-Waterloo, King's College London) for her assistance with sample analysis, and Ana Rodriguez-Mateos (Department of Nutritional Sciences, King's College London) for reading and commenting on the initial manuscript draft.

## **Financial Support**

CRESSIDA study was funded from the UK Food Standards Agency and Department of Health and by the National Institute for Health Research (NIHR) Clinical Research Facility at Guy's and St Thomas' NHS Foundation Trust and NIHR Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust and King's College London. M. L. C. A. received PhD studentship funding from the Mexican Secretariat of Public Education.

## **Conflict of Interest**

W.L.H. has current collaboration with Lucozade Ribena Suntory and DIANA Food SAS. T.A.B.S., D.P.R., J.D and M.L.C.A. reported no conflicts of interest.

## **Authorship**

T.A.B.S. was principal investigator of the CRESSIDA study. T.A.B.S. and W.L.H. devised the CRESSIDA study, D.P.R. and J.D. recruited subjects into the study and supported the dietary intervention in the CRESSIDA study. M.L.C.A. and W.L.H. conceived the idea for secondary analysis of polyphenols intake data, analysed data, performed statistical analysis. M.L.C.A. wrote drafts of the manuscripts, which were edited by W.L.H. The manuscript was read, commented on, and approved by all authors.

## References

1. Boeing H, Bechthold A, Bub A *et al.* (2012) Critical review: vegetables and fruit in the prevention of chronic diseases. *Eur J Nutr* **51**, 637-663.
2. Muraki I, Imamura F, Manson JE *et al.* (2013) Fruit consumption and risk of type 2 diabetes: results from three prospective longitudinal cohort studies. *BMJ* **347**, f5001.
3. Mursu J, Virtanen JK, Tuomainen TP *et al.* (2014) Intake of fruit, berries, and vegetables and risk of type 2 diabetes in Finnish men: the Kuopio Ischaemic Heart Disease Risk Factor Study. *Am J Clin Nutr* **99**, 328-333.
4. Buil-Cosiales P, Toledo E, Salas-Salvado J *et al.* (2016) Association between dietary fibre intake and fruit, vegetable or whole-grain consumption and the risk of CVD: results from the PREvencion con DIeta MEDiterranea (PREDIMED) trial. *Br J Nutr* **116**, 534-546.
5. Aune D, Keum N, Giovannucci E *et al.* (2016) Whole grain consumption and risk of cardiovascular disease, cancer, and all cause and cause specific mortality: systematic review and dose-response meta-analysis of prospective studies. *BMJ* **353**, i2716.
6. Ötles S, Ozgoz S (2014) Health effects of dietary fiber. *Acta Sci Pol Technol Aliment* **12**, 191-202.
7. Woodside JV, McCall D, McGartland C *et al.* (2005) Micronutrients: dietary intake v. supplement use. *Proc Nutr Soc* **64**, 543-553.
8. Rodriguez-Mateos A, Vauzour D, Krueger CG *et al.* (2014) Bioavailability, bioactivity and impact on health of dietary flavonoids and related compounds: an update. *Arch Toxicol* **88**, 1803-1853.
9. WHO (1990) *Diet, nutrition and the prevention of chronic diseases. Technical report series 797*. Geneva: World Health Organization
10. NHS (2005) 5 a day - Live well. <http://www.nhs.uk/livewell/5aday/Pages/5ADAYhome.aspx> (accessed 16/03 2016)
11. Bates B, Lennox A, Prentice A *et al.* (2014) National Diet and Nutrition Survey: Results from Years 1-4 (combined) of the Rolling Programme (2008/2009 – 2011/12) Executive summary [Health Do, editor]. London: Public Health England.
12. Mann KD, Pearce MS, McKeivith B *et al.* (2015) Low whole grain intake in the UK: results from the National Diet and Nutrition Survey rolling programme 2008-11. *Br J Nutr* **113**, 1643-1651.
13. Manach C, Scalbert A, Morand C *et al.* (2004) Polyphenols: food sources and bioavailability. *Am J Clin Nutr* **79**, 727-747.
14. Hooper L, Kroon PA, Rimm EB *et al.* (2008) Flavonoids, flavonoid-rich foods, and cardiovascular risk: a meta-analysis of randomized controlled trials. *Am J Clin Nutr* **88**, 38-50.
15. Jennings A, Welch AA, Fairweather-Tait SJ *et al.* (2012) Higher anthocyanin intake is associated with lower arterial stiffness and central blood pressure in women. *Am J Clin Nutr* **96**, 781-788.
16. McCullough ML, Peterson JJ, Patel R *et al.* (2012) Flavonoid intake and cardiovascular disease mortality in a prospective cohort of US adults. *Am J Clin Nutr* **95**, 454-464.
17. Cassidy A, Mukamal KJ, Liu L *et al.* (2013) High anthocyanin intake is associated with a reduced risk of myocardial infarction in young and middle-aged women. *Circulation* **127**, 188-196.
18. van Dam RM, Naidoo N, Landberg R (2013) Dietary flavonoids and the development of type 2 diabetes and cardiovascular diseases: review of recent findings. *Curr Opin Lipidol* **24**, 25-33.
19. Neuhouwer ML (2004) Dietary flavonoids and cancer risk: evidence from human population studies. *Nutr Cancer* **50**, 1-7.
20. Arts IC, Hollman PC (2005) Polyphenols and disease risk in epidemiologic studies. *Am J Clin Nutr* **81**, 317S-325S.
21. Fink BN, Steck SE, Wolff MS *et al.* (2007) Dietary flavonoid intake and breast cancer survival among women on Long Island. *Cancer Epidemiol Biomarkers Prev* **16**, 2285-2292.

22. Yuan JM (2011) Green tea and prevention of esophageal and lung cancers. *Mol Nutr Food Res* **55**, 886-904.
23. Wedick NM, Pan A, Cassidy A *et al.* (2012) Dietary flavonoid intakes and risk of type 2 diabetes in US men and women. *Am J Clin Nutr* **95**, 925-933.
24. Curtis PJ, Sampson M, Potter J *et al.* (2012) Chronic ingestion of flavan-3-ols and isoflavones improves insulin sensitivity and lipoprotein status and attenuates estimated 10-year CVD risk in medicated postmenopausal women with type 2 diabetes: a 1-year, double-blind, randomized, controlled trial. *Diabetes Care* **35**, 226-232.
25. Jennings A, Welch AA, Spector T *et al.* (2014) Intakes of anthocyanins and flavones are associated with biomarkers of insulin resistance and inflammation in women. *J Nutr* **144**, 202-208.
26. Jacques PF, Cassidy A, Rogers G *et al.* (2013) Higher dietary flavonol intake is associated with lower incidence of type 2 diabetes. *J Nutr* **143**, 1474-1480.
27. Zamora-Ros R, Knaze V, Lujan-Barroso L *et al.* (2011) Estimated dietary intakes of flavonols, flavanones and flavones in the European Prospective Investigation into Cancer and Nutrition (EPIC) 24 hour dietary recall cohort. *Br J Nutr* **106**, 1915-1925.
28. Zamora-Ros R, Knaze V, Lujan-Barroso L *et al.* (2011) Estimation of the intake of anthocyanidins and their food sources in the European Prospective Investigation into Cancer and Nutrition (EPIC) study. *Br J Nutr* **106**, 1090-1099.
29. Vogiatzoglou A, Mulligan AA, Luben RN *et al.* (2013) Assessment of the dietary intake of total flavan-3-ols, monomeric flavan-3-ols, proanthocyanidins and theaflavins in the European Union. *Br J Nutr*, **111**, 1463-1473.
30. Zamora-Ros R, Knaze V, Rothwell JA *et al.* (2015) Dietary polyphenol intake in Europe: the European Prospective Investigation into Cancer and Nutrition (EPIC) study. *Eur J Nutr* **55**, 1359-1375.
31. Yahya HM, Day A, Lawton C *et al.* (2015) Dietary intake of 20 polyphenol subclasses in a cohort of UK women. *Eur J Nutr* **55**, 1839-1847.
32. Pinto P, Santos CN (2017) Worldwide (poly)phenol intake: assessment methods and identified gaps. *Eur J Nutr* **56**, 1393-1408.
33. Naska A, Lagiou A, Lagiou P. Dietary assessment methods in epidemiological research: current state of the art and future prospects. *F1000Res*. Published online: 16 June 2017. doi: 10.12688/f1000research.10703.1
34. Somerset S, Papier K (2014) A food frequency questionnaire validated for estimating dietary flavonoid intake in an Australian population. *Nutr Cancer* **66**, 1200-1210.
35. Theodoratou E, Kyle J, Cetnarskyj R *et al.* (2007) Dietary flavonoids and the risk of colorectal cancer. *Cancer Epidemiol Biomarkers Prev* **16**, 684-693.
36. Zamora-Ros R, Touillaud M, Rothwell JA *et al.* (2014) Measuring exposure to the polyphenol metabolome in observational epidemiologic studies: current tools and applications and their limits. *Am J Clin Nutr* **100**, 11-26.
37. Neveu V, Perez-Jimenez J, Vos F *et al.* (2010) Phenol-Explorer: an online comprehensive database on polyphenol contents in foods. *Database (Oxford)* **2010**, bap024.
38. Rothwell JA, Urpi-Sarda M, Boto-Ordonez M *et al.* (2012) Phenol-Explorer 2.0: a major update of the Phenol-Explorer database integrating data on polyphenol metabolism and pharmacokinetics in humans and experimental animals. *Database (Oxford)* **2012**, bas031.
39. Rothwell JA, Perez-Jimenez J, Neveu V *et al.* (2013) Phenol-Explorer 3.0: a major update of the Phenol-Explorer database to incorporate data on the effects of food processing on polyphenol content. *Database* **2013**, bat070.
40. Nutrient Data Laboratory BHNRC, Agricultural Research Service (2004) USDA database for the proanthocyanidin content of selected foods. U.S: U.S. Department of Agriculture.

41. Bhagwat S, Haytowitz, D. B and Holden J. M. (2008) USDA database for the isoflavone content of selected foods. Release 2.0, September 2008 ed. U.S.: U.S. Department of Agriculture.
42. Bhagwat S, Haytowitz, D.B. Holden, J.M. (Ret) (2013) Database for the flavonoid content of selected foods, release 3.1. U.S.: USDA. Department of Agriculture, Agricultural Research Service.
43. Mennen LI, Sapinho D, Ito H *et al.* (2008) Urinary excretion of 13 dietary flavonoids and phenolic acids in free-living healthy subjects - variability and possible use as biomarkers of polyphenol intake. *Eur J Clin Nutr* **62**, 519-525.
44. Perez-Jimenez J, Hubert J, Hooper L *et al.* (2010) Urinary metabolites as biomarkers of polyphenol intake in humans: a systematic review. *Am J Clin Nutr* **92**, 801-809.
45. Nielsen S, Freese R, Kleemola P *et al.* (2002) Flavonoids in human urine as biomarker for intake of fruits and vegetables. *Cancer Epidemiol Biomarkers Prev* **11**, 459-466.
46. Krogholm KS, Haraldsdóttir J, Knuthsen P *et al.* (2004) Urinary Total Flavonoid Excretion but Not 4-Pyridoxic Acid or Potassium Can Be Used as a Biomarker for the Intake of Fruits and Vegetables. *J Nutr* **134**, 445-451.
47. Ito H, Gonthier M-P, Manach C *et al.* (2005) Polyphenol levels in human urine after intake of six different polyphenol-rich beverages. *Br J Nutr* **94**, 500-509.
48. Mullen W, Borges G, Lean ME *et al.* (2010) Identification of metabolites in human plasma and urine after consumption of a polyphenol-rich juice drink. *J Agric Food Chem* **58**, 2586-2595.
49. Mennen LI, Sapinho D, Ito H *et al.* (2006) Urinary flavonoids and phenolic acids as biomarkers of intake for polyphenol-rich foods. *Br J Nutr* **96**, 191-198.
50. Edmands WM, Ferrari P, Rothwell JA *et al.* (2015) Polyphenol metabolome in human urine and its association with intake of polyphenol-rich foods across European countries. *Am J Clin Nutr* **102**, 905-913.
51. Spencer JP, Abd El Mohsen MM, Minihaue AM *et al.* (2008) Biomarkers of the intake of dietary polyphenols: strengths, limitations and application in nutrition research. *Br J Nutr* **99**, 12-22.
52. Williamson G, Holst B (2008) Dietary reference intake (DRI) value for dietary polyphenols: are we heading in the right direction? *Br J Nutr* **99** Suppl 3, S55-58.
53. Lupton JR, Atkinson SA, Chang N *et al.* (2014) Exploring the benefits and challenges of establishing a DRI-like process for bioactives. *Eur J Nutr* **53** Suppl 1, 1-9.
54. Williamson G (2017) The role of polyphenols in modern nutrition. *Nutr Bull* **42**, 226-235.
55. Perez-Jimenez J, Neveu V, Vos F *et al.* (2010) Systematic analysis of the content of 502 polyphenols in 452 foods and beverages: an application of the phenol-explorer database. *J Agric Food Chem* **58**, 4959-4969.
56. Reidlinger DP, Darzi J, Hall WL *et al.* (2015) How effective are current dietary guidelines for cardiovascular disease prevention in healthy middle-aged and older men and women? A randomized controlled trial. *Am J Clin Nutr* **101**, 922-930.
57. Reidlinger DP (2015) Evidence, benefits and barriers to achieving an integrated cardioprotective dietary pattern. PhD Thesis, King's College London.
58. Finglas PM, Roe, M. A., Pinchen, H. M., Berry, B., Church, S. M., Dodhia, S.K., Farron-Wilson, M. and Swan, G. (2015) *McCance and Widdowson's The composition of foods*. Seventh Summary Edition ed. Cambridge: Royal Society of Chemistry
59. Campbell J, Foskett, D., Rippington, N., Paskins, P. (2012) *Practical cookery*. 12th Edition ed. UK: Hodder Education.
60. Worldwide B (2009) BBC good food. <http://www.bbcgoodfood.com/>
61. Bingham SA, C. G, Welch A *et al.* (1994) Comparison of dietary assessment methods in nutritional epidemiology: weighed records v. 24h recalls, food frequency questionnaires and estimated diet records. *B J Nut* **72**, 619-642.

62. Achaintre D, Buleté A, Cren-Olivé C *et al.* (2016) Differential isotope labeling of 38 dietary polyphenols and their quantification in urine by liquid chromatography electrospray ionization tandem mass spectrometry. *Anal Chem* **88**, 2637-2644.
63. Bingham S, Cummings JH (1983) The use of 4-aminobenzoic acid as a marker to validate the completeness of 24 h urine collections in man. *Clin Sci* **64**, 629-635.
64. Rhee JJ, Sampson L, Cho E *et al.* (2014) Comparison of methods to account for implausible reporting of energy intake in epidemiologic studies. *Am J Epidemiol* **181**, 225-233.
65. Tetens I, Turrini A, Tapanainen H *et al.* Dietary intake and main sources of plant lignans in five European countries. *Food Nutr Res*. Published online: 11 June 2013. doi: 10.3402/fnr.v57i0.19805.
66. Mulligan AA, Kuhnle GG, Lentjes MA *et al.* (2013) Intakes and sources of isoflavones, lignans, enterolignans, coumestrol and soya-containing foods in the Norfolk arm of the European Prospective Investigation into Cancer and Nutrition (EPIC-Norfolk), from 7 d food diaries, using a newly updated database. *Public Health Nutr* **16**, 1454-1462.
67. Knaze V, Zamora-Ros R, Lujan-Barroso L *et al.* (2012) Intake estimation of total and individual flavan-3-ols, proanthocyanidins and theaflavins, their food sources and determinants in the European Prospective Investigation into Cancer and Nutrition (EPIC) study. *Br J Nutr* **108**, 1095-1108.
68. Bingham SA GC, Welch A, Day K, Cassidy A, Khaw K T, Sneyd MJ, Key TJA, Roe L and Day NE. (1994) Comparison of dietary assessment methods in nutritional epidemiology: weighed records v. 24h recalls, food frequency questionnaires and estimated diet records. *B J Nut* **72**, 619-642.
69. Zamora-Ros R, Rabassa M, Llorach R *et al.* (2012) Application of Dietary Phenolic Biomarkers in Epidemiology: Past, Present, and Future. *J Agric Food Chem* **60**, 6648–6657.
70. Chen Z, Zheng S, Li L *et al.* (2014) Metabolism of flavonoids in human: A comprehensive review. *Current Drug Metabolism* **15**, 48-61.
71. Saha S, Hollands W, Needs PW, Ostertag LM, de Roos B, Duthie GG and Kroon PA (2012) Human O-sulfated metabolites of (–)-epicatechin and methyl-(–)-epicatechin are poor substrates for commercial aryl-sulfatases: Implications for studies concerned with quantifying epicatechin bioavailability. *Pharmacol Res* **65**, 592-602.
72. Henning SM, Wang P, Abgaryan N *et al.* (2013) Phenolic acid concentrations in plasma and urine from men consuming green or black tea and potential chemopreventive properties for colon cancer. *Mol Nutr Food Res* **57**, 483-493.
73. Crozier A, Clifford MN, Del rio D (2012) Bioavailability of dietary monomeric and polymeric flavan-3-ols. In *Flavonoids and related compounds*, pp. 45-78 [Spencer JPE and Crozier A, editors]. Boca Raton, FL.: CRC Press.
74. Nurmi T, Mursu J, Heinonen M *et al.* (2009) Metabolism of berry anthocyanins to phenolic acids in humans. *J Agric Food Chem* **57**, 2274-2281.
75. de Ferrars RM, Cassidy A, Curtis P *et al.* (2013) Phenolic metabolites of anthocyanins following a dietary intervention study in post-menopausal women. *Mol Nutr Food Res* **58**, 490-502.
76. Olthof MR, Hollman PC, Buijsman MN *et al.* (2003) Chlorogenic acid, quercetin-3-rutinoside and black tea phenols are extensively metabolized in humans. *J Nutr* **133**, 1806-1814.
77. Rios LY, Gonthier MP, Remesy C *et al.* (2003) Chocolate intake increases urinary excretion of polyphenol-derived phenolic acids in healthy human subjects. *Am J Clin Nutr* **77**, 912-918.
78. Pounis G, Di Castelnuovo A, Bonaccio M *et al.* (2016) Flavonoid and lignan intake in a Mediterranean population: proposal for a holistic approach in polyphenol dietary analysis, the Moli-sani Study. *Eur J Clin Nutr* **70**, 338-345.
79. Valls-Pedret C, Lamuela-Raventos RM, Medina-Remon A *et al.* (2012) Polyphenol-rich foods in the Mediterranean diet are associated with better cognitive function in elderly subjects at high cardiovascular risk. *J Alzheimers Dis* **29**, 773-782.

80. Shi Y, Williamson G (2015) Comparison of the urinary excretion of quercetin glycosides from red onion and aglycone from dietary supplements in healthy subjects: a randomized, single-blinded, cross-over study. *Food Funct* **6**, 1443-1448.
81. Chong MF, George TW, Alimbetov D *et al.* (2012) Impact of the quantity and flavonoid content of fruits and vegetables on markers of intake in adults with an increased risk of cardiovascular disease: the FLAVURS trial. *Eur J Nutr* **52**, 361-78
82. Brevik A, Rasmussen SE, Dreven CA *et al.* (2004) Urinary excretion of flavonoids reflects even small changes in the dietary intake of fruits and vegetables. *Cancer Epidemiol Biomarkers Prev* **13**, 843-849.
83. Vetrani C, Rivellese AA, Annuzzi G *et al.* (2014) Phenolic metabolites as compliance biomarker for polyphenol intake in a randomized controlled human intervention. *Food Res Int* **63**, 233-238.
84. Garcia-Conesa MT, Chambers K, Combet E *et al.* Meta-analysis of the effects of foods and derived products containing ellagitannins and anthocyanins on cardiometabolic biomarkers: analysis of factors influencing variability of the individual responses. *Int J Mol Sci*. Published online: 19 February 2018. doi: 10.3390/ijms19030694.
85. Rienks J, Barbaresko J, Nothlings U (2017) Association of isoflavone biomarkers with risk of chronic disease and mortality: a systematic review and meta-analysis of observational studies. *Nutr Rev* **75**, 616-641.
86. Onakpoya IJ, Spencer EA, Thompson MJ *et al.* (2015) The effect of chlorogenic acid on blood pressure: a systematic review and meta-analysis of randomized clinical trials. *J Hum Hypertens* **29**, 77-81.
87. Testai L, Calderone V. Nutraceutical Value of Citrus Flavanones and Their Implications in Cardiovascular Disease. *Nutrients*. Published online: 16 May 2017 May. doi: 10.3390/nu9050502.
88. Törrönen R, Kolehmainen M, Sarkkinen E *et al.* (2012) Postprandial glucose, insulin, and free fatty acid responses to sucrose consumed with blackcurrants and lingonberries in healthy women. *Am J Clin Nutr* **96**, 527-533.
89. Törrönen R, Sarkkinen E, Niskanen T *et al.* (2012) Postprandial glucose, insulin and glucagon-like peptide 1 responses to sucrose ingested with berries in healthy subjects. *Br J Nutr* **107**, 1445-1451.
90. Törrönen R, Sarkkinen E, Tapola N *et al.* (2010) Berries modify the postprandial plasma glucose response to sucrose in healthy subjects. *Br J Nutr* **103**, 1094-1097.
91. Castro-Acosta ML, Smith L, Miller RJ *et al.* (2016) Drinks containing anthocyanin-rich blackcurrant extract decrease postprandial blood glucose, insulin and incretin concentrations. *J Nutr Biochem* **38**, 154-161.
92. Hartley L, Flowers N, Holmes J *et al.* Green and black tea for the primary prevention of cardiovascular disease. *Cochrane Database Syst Rev*. Published online: 18 June 2013 doi: 10.1002/14651858.CD009934.pub2.
93. Tajik N, Tajik M, Mack I *et al.* (2017) The potential effects of chlorogenic acid, the main phenolic components in coffee, on health: a comprehensive review of the literature. *Eur J Nutr* **56**, 2215-2244.
94. Dragsted LO, Gao Q, Pratico G *et al.* Dietary and health biomarkers-time for an update. *Genes Nutr*. Published online: 29 September 2017. doi: 10.1186/s12263-017-0578-y.
95. Noh H, Freisling H, Assi N *et al.* Identification of urinary polyphenol metabolite patterns associated with polyphenol-rich food intake in adults from four European countries. *Nutrients*. Published online: 25 July 2017. doi: 10.3390/nu9080796.
96. Woodside JV, Draper J, Lloyd A *et al.* (2017) Use of biomarkers to assess fruit and vegetable intake. *Proc Nutr Soc* **76**, 308-315.
97. Brouwer-Brolsma EM, Brennan L, Dreven CA *et al.* (2017) Combining traditional dietary assessment methods with novel metabolomics techniques: present efforts by the Food Biomarker Alliance. *Proc Nutr Soc* **76**, 619-627.



98. Kuhnle GG (2012) Nutritional biomarkers for objective dietary assessment. *J Sci Food Agric* **92**, 1145-1149.
99. Cassidy A, O'Reilly EJ, Kay C *et al.* (2011) Habitual intake of flavonoid subclasses and incident hypertension in adults. *Am J Clin Nutr* **93**, 338-347.
100. McKeown NM, Day NE, Welch AA *et al.* (2001) Use of biological markers to validate self-reported dietary intake in a random sample of the European Prospective Investigation into Cancer United Kingdom Norfolk cohort. *Am J Clin Nutr* **74**, 188-196.
101. Ranka S, Gee JM, Biro L *et al.* (2008) Development of a food frequency questionnaire for the assessment of quercetin and naringenin intake. *Eur J Clin Nutr* **62**, 1131-1138.
102. Zhang Y, Cao J, Chen W *et al.* (2010) Reproducibility and relative validity of a food frequency questionnaire to assess intake of dietary flavonol and flavone in Chinese university campus population. *Nutr Res* **30**, 520-526.
103. Mullie P, Clarys P, Deriemaeker P *et al.* (2007) Estimation of daily human intake of food flavonoids. *Plant Foods Hum Nutr* **62**, 93-98.
104. Cheynier V, Tomas-Barberan FA, Yoshida K (2015) Polyphenols: From Plants to a Variety of Food and Nonfood Uses. *J Agric Food Chem* **63**, 7589-7594.
105. Ding Y, Peng M, Zhang T *et al.* (2013) Quantification of conjugated metabolites of drugs in biological matrices after the hydrolysis with  $\beta$ -glucuronidase and sulfatase: a review of bio-analytical methods. *Biomed Chromatogr* **27**, 1280-1295.
106. Burkholder-Cooley NM, Rajaram SS, Haddad EH *et al.* (2017) Validating polyphenol intake estimates from a food-frequency questionnaire by using repeated 24-h dietary recalls and a unique method-of-triads approach with 2 biomarkers. *Am J Clin Nutr* **105**, 685-694.

## **Figure legends**

**Figure 1** Flow of participants through the CRESSIDA study

**Figure 2** Main polyphenol food sources in control and dietary guidelines groups, at endpoint, of the CRESSIDA study as estimated by four-day food diary

**Figure 3** Percentage changes from baseline in main food group sources of polyphenols in dietary guidelines group as estimated by four-day food diary

**Table 1** Baseline characteristics of the CRESSIDA study population by randomised group

Characteristic	Control (n=82)		Dietary Guidelines (n=79)		Difference between groups ( <i>P</i> * <i>value</i> )
Gender (n)					
Female	50		47		
Male	32		32		
	Mean	SD	Mean	SD	
Age (years)	52	8.0	53	8.0	0.717
BMI (kg/m <sup>2</sup> )	26.8	3.9	25.4	3.7	0.019
Waist circumference (cm)					
Female/male	91.9/98.2	10.0/12.2	87.1/97.3	12.2/9.3	0.093
SBP (mmHg)	122.0	11.5	121.6	14.3	0.848
DBP (mmHg)	73.9	7.1	73.8	8.0	0.930
Glucose (mmol/L)	5.0	0.4	5.3	0.5	0.990
Total cholesterol (mmol/L)	5.3	0.9	5.4	1.1	0.983
TC:HDL cholesterol ratio	3.6	1.0	3.5	1.0	0.262

BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure.

\*Statistical comparisons between groups at baseline by independent T-test or independent samples Mann-Whitney U-test.

**Table 2** Daily intake of polyphenols estimated by four-day food diary and adjusted for energy intake following dietary guidelines and control diets.

Polyphenols (mg/d/10 MJ) Classes and <i>subclasses</i>	Baseline (n=161)		Endpoint (n=161)				Differences between groups at endpoint ( <i>P</i> <sup>‡</sup> )
			Control (n=82)		Dietary Guidelines (n=79)		
Total polyphenols*	1202 <sup>†</sup>	(1093, 1323)	1084	(980, 1197)	1279	(1158, 1412)	0.023 <sup>§</sup>
Flavonoids	840 <sup>†</sup>	(536, 1263)	802	(456, 1109)	882	(610, 1401)	0.014
<i>Anthocyanidins</i>	55	(14, 130)	30	(14, 108)	76	(25, 144)	0.032
<i>Dihydrochalcones</i>	1.6	(0.0, 3.8)	2	(0, 4)	2	(1, 6)	0.297
<i>Flavanols</i>	349	(114, 526)	311	(104, 498)	394	(128, 665)	0.159
<i>Proanthocyanidins</i>	231 <sup>†</sup>	(117, 339)	195	(115, 324)	249	(169, 381)	0.020
<i>Flavanones</i>	31	(4, 64)	24	(2, 58)	33	(8, 84)	0.096
<i>Flavones</i>	6	(3, 12)	5	(1, 12)	5	(2, 12)	0.603
<i>Flavonols</i>	80	(51, 112)	80	(50, 105)	83	(58, 127)	0.125
<i>Isoflavones</i>	0.3	(0.1, 9.9)	0.1	(0.1, 0.4)	0.5	(0.1, 65.0)	0.002
Lignans	0.5	(0.2, 5.0)	0.3	(0.1, 0.4)	0.5	(0.2, 6.0)	<0.001
Phenolic acids	289	(185, 569)	221	(132, 493)	344	(209, 522)	0.012
<i>Hydroxybenzoic acids</i>	90	(35, 143)	75	(32, 125)	100	(55, 170)	0.067
<i>Hydroxycinnamic acids</i>	149	(88, 450)	119	(59, 378)	172	(99, 444)	0.018
Stilbenes	0.09	(0.02, 0.71)	0.09	(0.02, 0.94)	0.26	(0.07, 1.25)	0.042
Other polyphenols	21	(11, 41)	13	(6, 30)	37	(20, 58)	<0.001

All values are medians (with lower and upper limits of IQR), except where \* denotes values are geometric means (95% CI), adjusted for baseline and gender at endpoint.

†Statistically significant differences between groups at baseline by independent T-test (total polyphenols) or independent samples Mann-Whitney U-test (all other comparisons); intakes were higher in the DG group than the control group,  $P<0.05$ .

‡Statistical comparisons between groups at endpoint by Mann-Whitney U-test, except where § denotes statistical comparisons between groups at endpoint adjusted for baseline and gender by ANCOVA.

**Table 3** Median daily intake of flavonoids estimated by food frequency questionnaire in dietary guidelines and control groups

Flavonoids (mg/d) Class and <i>subclasses</i>	Baseline (n=161)		Endpoint (n=161)				Differences between groups at endpoint ( <i>P</i> *)
			Control (82)		Dietary Guidelines (79)		
Total flavonoids	661	(417, 867)	539	(350, 862)	715	(571, 906)	0.005
<i>Anthocyanidins</i>	19	(10, 35)	16	(9, 31)	32	(19, 50)	<0.001
<i>Flavones</i>	2.4	(1.6, 3.5)	2.7	(1.5, 3.7)	3.2	(2.0, 4.4)	0.013
<i>Flavonols</i>	35	(23, 47)	39	(21, 54)	39	(31, 50)	0.199
<i>Flavanols</i>	90	(40, 143)	85	(35, 146)	97	(80, 143)	0.048
<i>Flavanones</i>	23	(8, 42)	25	(10, 43)	27	(8, 49)	0.352
<i>Proanthocyanidins</i>	265	(184, 379)	232	(184, 371)	290	(229, 385)	0.021

Values are medians (lower and upper limits of the IQR). There were no significant differences between groups at baseline.

\* Statistical comparisons between groups at endpoint by Mann-Whitney U-test.

**Table 4** Comparison of urinary phenolic excretion method and polyphenol intake estimated by 4-day food diary in their ability to discriminate between subsample populations either adhering to dietary guidelines or a control diet.

Polyphenols	Urinary excretion <sup>†</sup> (μmol/d)					Dietary intake (mg/d/10 MJ)				
	Control (n=46)		Dietary Guidelines (n=45)		Difference between groups ( <i>P</i> )	Control (n=46)		Dietary Guidelines (n=45)		Difference between groups ( <i>P</i> )
Phloretin <sup>‡</sup>	0.09	(0.01, 0.28)	0.40*	(0.18, 0.97)	<0.001	1.2	(0.0, 3.5)	3.1*	(1.1, 5.9)	0.006
Epicatechin	0.45	(0.24, 0.60)	0.44	(0.21, 0.70)	0.943	30	(15, 50)	40*	(29, 67)	0.013
Eriodictyol	0.25	(0.09, 0.43)	0.51*	(0.20, 0.86)	0.011	0.00	(0.00, 0.02)	0.02*	(0.01, 0.06)	<0.001
Hesperetin	0.20	(0.00, 2.03)	1.38*	(0.09, 3.51)	0.010	8	(1, 23)	17*	(4, 35)	0.019
Luteolin	1.8	(0.9, 2.9)	2.7*	(1.5, 3.7)	0.017	0.5	(0.3, -8)	1.5*	(0.9, 3.3)	<0.001
Quercetin	4.7	(1.5, 7.4)	6.6	(2.4, 12.8)	0.105	50	(27, 66)	59*	(41, 85)	0.028
Daidzein	0.02	(0.00, 0.40)	0.11	(0.00, 0.80)	0.074	0.04	(0.04, 0.10)	3.5*	(0.02, 21.2)	0.006
Gallic acid	0.7	(0.2, 2.0)	1.4	(0.5, 3.6)	0.053	62	(18, 107)	90	(23, 153)	0.134
Vanillic acid	1.4	(0.2, 2.6)	3.0*	(0.9, 5.9)	0.023	0.2	(0.1, 0.2)	0.4*	(0.2, 0.6)	<0.001
Lignan/ Enterolactone <sup>§</sup>	5.7	(2.1, 10.6)	10.1*	(3.8, 22.6)	0.010	0.2	(0.1, 0.4)	0.6*	(0.3, 6.1)	<0.001

Excretion of polyphenols in 24 h urine and dietary intake of the direct dietary precursor estimated by 4-DFD at endpoint in a subsample of the CRESSIDA study population selected for the lowest (control group) and highest (dietary guidelines group) fruit and vegetables intake using self-reported data from food frequency questionnaires (n=91).


All values are median (lower and upper limits of IQR). All between group comparisons were carried out by independent samples Mann-Whitney U Test.

<sup>†</sup> Values adjusted for 24 h urine volume.

<sup>‡</sup> Sum of phloretin and phlorizin

<sup>§</sup> Intake of lignans was used as direct dietary precursor of enterolactone.


\*Statistically significant differences between groups at endpoint, *P*<0.05.



Control diet

Dietary guideline diet


Total polyphenols




Total polyphenols




Flavonoids



Flavonoids



Flavanols



Flavanols

